

Erratum

Statistics review 3: Hypothesis testing and *P* values

Elise Whitley¹ and Jonathan Ball²

¹Lecturer in Medical Statistics, University of Bristol, UK

²Lecturer in Intensive Care Medicine, St George's Hospital Medical School, London, UK

Correspondence: Editorial Office, *Critical Care*, editorial@ccforum.com

Published online: 13 December 2002

Critical Care 2003, **7**:15 (DOI 10.1186/cc1868)

This article is online at <http://ccforum.com/content/7/1/15>

© 2003 BioMed Central Ltd (Print ISSN 1364-8535; Online ISSN 1466-609X)

An erratum to *Statistics review 3: Hypothesis testing and P values*, by Elise Whitley and Jonathan Ball.

In the original version of this article [1], the words 'if the null hypothesis is true' were omitted from the first sentence of the first key message. The key message should have read as follows:

'A *P* value is the probability that an observed effect is simply due to chance, if the Null hypothesis is true; it therefore provides a measure of the strength of an association. A *P* value does not provide any measure of the size of an effect, and cannot be used in isolation to inform clinical judgement.'

References

1. Whitley E, Ball J: **Statistics review 3: Hypothesis testing and P values**. *Crit Care* 2002, **6**:222-225.